



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Preterm birth prevention-Time to PROGRESS beyond progesterone

Citation for published version:

Norman, JE & Bennett, P 2017, 'Preterm birth prevention-Time to PROGRESS beyond progesterone', *PLoS Medicine*, vol. 14, no. 9, pp. e1002391. <https://doi.org/10.1371/journal.pmed.1002391>

Digital Object Identifier (DOI):

[10.1371/journal.pmed.1002391](https://doi.org/10.1371/journal.pmed.1002391)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

PLoS Medicine

Publisher Rights Statement:

This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.




PERSPECTIVE

Preterm birth prevention—Time to PROGRESS beyond progesterone

Jane E. Norman^{1*}, Phillip Bennett²

1 MRC Centre for Reproductive Health, University of Edinburgh, Edinburgh, United Kingdom, **2** Institute for Reproductive and Developmental Biology, Imperial College London, London, United Kingdom

* jane.norman@ed.ac.uk



OPEN ACCESS

Citation: Norman JE, Bennett P (2017) Preterm birth prevention—Time to PROGRESS beyond progesterone. *PLoS Med* 14(9): e1002391. <https://doi.org/10.1371/journal.pmed.1002391>

Published: September 26, 2017

Copyright: © 2017 Norman, Bennett. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: JEN and PB are supported by the charity Tommy's. The funders had no role in decision to publish, or preparation of the manuscript.

Competing interests: JEN and PB were co-authors and co-grant holders of the OPPTIMUM study. OPPTIMUM was funded by the Efficacy and Mechanism Evaluation (EME) programme, a Medical Research Council (MRC) and National Institute of Health Research (NIHR) partnership, award number G0700452, revised to 09/800/27. The EME programme is funded by the MRC and NIHR, with contributions from the Chief Scientist Office in Scotland and National Institute for Social Care and Research in Wales. PB is a consultant to and holds shares in ObsEva, which develops drugs for the prevention and/or delay of preterm birth.

Abbreviations: FDA, United States Food and Drug Administration.

In this week's *PLOS Medicine*, Crowther and colleagues publish the results of the PROGRESS Study, which examined the effect of maternal vaginal progesterone on the risk of respiratory distress syndrome in the baby [1]. Women at high risk of preterm birth because of a previous spontaneous preterm birth were recruited to this study, which involved self-administering 100 mg progesterone or a placebo vaginally from 20 to 34 weeks of pregnancy. No differences were found in the rate of the primary outcome (incidence of respiratory distress syndrome and severity of respiratory disease) between the progesterone and the placebo groups (incidence of respiratory distress syndrome in 42/402 [10.5%] and 41/388 [10.6%], respectively [RR 0.98, 95% CI 0.64–1.49]), nor were there differences in any of the other secondary outcomes. The study, a double-masked randomised placebo-controlled trial, was well conducted. About 40% of women who were approached agreed to participate. The primary outcome was available for all participating women, and around 65% of women were still taking study treatment by 34 weeks gestation. Rates of preterm birth were 36% and 17% before 37 and 34 weeks, respectively; hence, the recruitment strategy effectively identified a group at high risk. Although the study was terminated earlier than planned (after recruitment of 787 women rather than the 984 initially intended), and the incidence of the primary outcome was lower than anticipated in the placebo group (10.6% rather than 15%), it is unlikely that these issues have materially affected the conclusions of the study.

Although vaginal progesterone is not licensed for prevention of preterm birth, it has been widely adopted by clinicians around the world for this purpose. Progesterone for preterm birth prevention in high-risk women is endorsed by expert guideline groups [2], including the National Institute for Clinical Excellence (UK) [3] and the FIGO Working Group on Best Practice in Maternal-Fetal Medicine [4]. The PROGRESS Study is the third large randomized trial (the others being our own OPPTIMUM study [5] and the study of O'Brien et al. [6]) to show a lack of efficacy of vaginal progesterone. Specifically, the PROGRESS trial showed no difference between the 2 groups in gestation of delivery, preterm birth before 37 weeks, or preterm birth before 34 weeks. We believe it is now time for a comprehensive review of the data on progesterone to identify risks and benefits in women at risk of preterm birth for different reasons.

Ostensibly, there is biological plausibility behind the administration of progesterone for prevention of preterm birth. Progesterone can inhibit contractions of the myometrium in vitro [7], and antiprogesterones are used as abortifacients. However, progesterone levels are high during pregnancy, far above receptor saturation. There is no evidence that women delivering preterm have lower progesterone levels and administration of progesterone vaginally

Provenance: Commissioned; not externally peer reviewed.

does not increase circulating concentrations. Therefore, the mechanism by which a modest additional amount of progesterone could achieve a therapeutic effect is unclear.

In evaluating progesterone to prevent preterm birth, the Cochrane analysis groups evidence by maternal risk factor for preterm birth rather than by route of administration or type of progesterone [8], even though vaginal natural progesterone and intramuscular 15 dihydroprogesterone caproate are very different therapies. The Cochrane analysis suggests that, in women with a previous preterm birth, progestogens reduce perinatal mortality and rates of preterm birth before 34 weeks, low birthweight, and neonatal death. Inclusion of women from OPPTIMUM and PROGRESS will double the existing sample size; a major impact on the treatment effect and confidence intervals is to be expected. In women at risk of preterm birth because of a short cervix, Cochrane suggests that progesterone reduces rates of preterm birth before 34 weeks but has no impact on perinatal mortality, low birthweight, or neonatal death [8]. A single randomized study and an individual patient data meta-analysis have shown that progesterone in women with a short cervix reduces a composite adverse neonatal outcome [9,10], although the individual study was insufficient to persuade the United States Food and Drug Administration (FDA) to license vaginal progesterone. Importantly, the FDA considered that a more robust statistical approach failed to result in a statistically significant effect of progesterone [11]. To address this complexity, the Patient-Centered Outcomes Research Initiative (PCORI) has initiated an individual patient data meta-analysis that will be conducted by an independent data centre. It is hoped that this will address the uncertainty about the efficacy of progesterone to prevent preterm birth in women with various risk factors.

Even if progesterone is found to reduce preterm birth rates in women with either a previous preterm birth or a short cervix, it is increasingly evident that alternatives are needed because the impact on preterm birth rates overall are small. Modelling of the impact of progesterone to prevent preterm birth in women with a previous preterm birth (but without a short cervix), and assuming a 20% reduction in preterm birth before 37 weeks, suggests an absolute reduction of 0.01% in preterm birth rates [12]. It has been claimed that routine cervical length screening of the entire US pregnant population, and offering vaginal progesterone to all women with a cervical length of 10–20 mm, would be cost effective [13]. However, in this scenario preterm birth would be prevented in only 913 women. This represents an absolute risk reduction of 0.02%, given that in 2015 there were 382,786 preterm births in the US [14].

There has been huge renewed excitement amongst clinicians and the pharmaceutical industry for the use of progestogens to prevent preterm birth in the 21st century. A PubMed search shows 792 publications on this topic since 2000. The PROGRESS Study adds to a body of evidence suggesting that the role of progesterone may be more limited than previously thought. Preterm birth remains the single biggest cause of neonatal mortality and morbidity worldwide, and effective therapeutic interventions are desperately needed. It is time to start exploring alternatives, and to progress beyond progesterone for the prevention of preterm birth.

References

1. Crowther CA, Ashwood P, McPhee AJ, Flenady V, Tran T, Dodd JM, et al. Vaginal progesterone pessaries for pregnant women with a previous preterm birth to prevent neonatal respiratory distress syndrome (the PROGRESS Study): A multicentre, randomised, placebo-controlled trial. *PLoS Med.* 2017; 14(9): e1002390. <https://doi.org/10.1371/journal.pmed.1002390>
2. Society for Maternal-Fetal Medicine Publications Committee, with assistance of Vincenzo Berghella. Progesterone and preterm birth prevention: translating clinical trials data into clinical practice. *Am J Obstet Gynecol.* 2012; 206(5):376–86. Epub 2012/05/01. <https://doi.org/10.1016/j.ajog.2012.03.010> PMID: 22542113.

3. National Institute for Health and Care Excellence. Preterm labour and birth. NICE guideline [NG25]. 2015. Available from: <https://www.nice.org.uk/guidance/ng25>.
4. FIGO Working Group on Best Practice in Maternal-Fetal Medicine. Best practice in maternal-fetal medicine. *Int J Gynecol Obstet*. 2015; 128(1):80–2. <https://doi.org/10.1016/j.ijgo.2014.10.011> PMID: 25481030
5. Norman JE, Marlow N, Messow CM, Shennan A, Bennett PR, Thornton S, et al. Vaginal progesterone prophylaxis for preterm birth (the OPPTIMUM study): a multicentre, randomised, double-blind trial. *Lancet*. 2016; 387:2106–16. [https://doi.org/10.1016/S0140-6736\(16\)00350-0](https://doi.org/10.1016/S0140-6736(16)00350-0) PMID: 26921136.
6. O'Brien JM, Adair CD, Lewis DF, Hall DR, DeFranco EA, Fusey S, et al. Progesterone vaginal gel for the reduction of recurrent preterm birth: primary results from a randomized, double-blind, placebo-controlled trial. *Ultrasound Obstet Gynecol*. 2007; 30:687–96. <https://doi.org/10.1002/uog.5158> PMID: 17899572
7. Anderson L, Martin W, Higgins C, Nelson S, Norman J. The effect of progesterone on myometrial contractility, potassium channels, and tocolytic efficacy. *Reprod Sci*. 2009; 16:1052–61. <https://doi.org/10.1177/1933719109340926> PMID: 19602723.
8. Dodd JM, Jones L, Flenady V, Cincotta R, Crowther CA. Prenatal administration of progesterone for preventing preterm birth in women considered to be at risk of preterm birth. *Cochrane Database Syst Rev*. 2013; 7:CD004947. <https://doi.org/10.1002/14651858.CD004947.pub3> PMID: 23903965.
9. Romero R, Nicolaides K, Conde-Agudelo A, Tabor A, O'Brien JM, Cetingoz E, et al. Vaginal progesterone in women with an asymptomatic sonographic short cervix in the midtrimester decreases preterm delivery and neonatal morbidity: a systematic review and metaanalysis of individual patient data. *Am J Obstet Gynecol*. 2012; 206(2):124 e1–19. Epub 2012/01/31. <https://doi.org/10.1016/j.ajog.2011.12.003> PMID: 22284156; PMCID: PMC3437773.
10. Hassan SS, Romero R, Vidyadhari D, Fusey S, Baxter J, Khandelwal M, et al. Vaginal progesterone reduces the rate of preterm birth in women with a sonographic short cervix: a multicenter, randomized, double-blind, placebo-controlled trial. *Ultrasound Obstet Gynecol*. 2011. Epub 2011/04/08. <https://doi.org/10.1002/uog.9017> PMID: 21472815.
11. Division of Reproductive and Urologic Products Office of New Drugs Center for Drug Evaluation and Research Food and Drug Administration. Background document for meeting of Advisory Committee for Reproductive Health Drugs. 2011. Available from: <https://wayback.archive-it.org/7993/20170405210910/https://www.fda.gov/downloads/AdvisoryCommittees/CommitteesMeetingMaterials/Drugs/ReproductiveHealthDrugsAdvisoryCommittee/UCM287987.pdf>
12. Chang HH, Larson J, Blencowe H, Spong CY, Howson CP, Cairns-Smith S, et al. Preventing preterm births: analysis of trends and potential reductions with interventions in 39 countries with very high human development index. *Lancet*. 2012. Epub 2012/11/20. [https://doi.org/10.1016/S0140-6736\(12\)61856-X](https://doi.org/10.1016/S0140-6736(12)61856-X) PMID: 23158883.
13. Einerson BD, Grobman WA, Miller ES. Cost-effectiveness of risk-based screening for cervical length to prevent preterm birth. *Am J Obstet Gynecol*. 2016; 215(1):100 e1–7. <https://doi.org/10.1016/j.ajog.2016.01.192> PMID: 26880732.
14. Martin JA, Hamilton BE, Osterman MJ, Driscoll AK, Mathews TJ. Births: final data for 2015. *Natl Vital Stat Rep*. 2017; 66(1):1. PMID: 28135188.